




## Lyofast Y 470 E

Description Application	<b>Lyofast Y 470 E</b> consists of specifically selected strains of <i>Streptococcus</i> <i>thermophilus</i> and <i>Lactobacillus delbrueckii</i> ssp. <i>bulgaricus</i> both producing EPS. Lyofast Y 470 E ensures a uniform and controlled production of traditional, aromatic, viscous yoghurt used for drinkable yoghurt, set and stirred yoghurt. Sprinkle the culture powder directly into process milk under aseptic conditions ensuring				
	that the culture is well dispersed by gentle stirring. The following may be used as inoculation guidelines:				
	Product	UC/100 I	Product	UC/100 I	
	Yoghurt, short set	2.0-4.0	Yoghurt, long set	0.5-1.0	
Rotation	The recommended rotations are Lyofast Y 472 E/Y 476 E.				
Acidification information	Standardised laboratory acidification test is conducted in milk powder, reconstituted at 10%, at defined temperature. Acidification profile: inoculation level corresponding to 1 UC per 100 litres milk.				

Standard activity: expressed as temperature/time/pH relations: 43°C/6 hours/pH 4.5 ± 0.15.



## Culture information

Data are obtained under standardised laboratory conditions, and consequently, should be considered as guidelines.

Optimal temperature for growth	43 °C	Urease activity	+
Acidification capability	pH 3.8	Texture formation	6 ± 1 sec/g
Aroma formation for yoghurt	++	Post-acidification	Δ pH 0.5

Storage

Unopened pouches should be kept below -17°C.

The freeze-dried culture is packed in waterproof and airproof aluminium pouches. The Package data packaging material is food grade. Lyofast Y 470 E is available in 10 and 50 UC. M91Y470E/4/UK/0 Issue: 04/02/2009 Review: 4 of 19/08/2014 1/2





## Lyofast Y 470 E

Shelf life

18 months when stored below -17℃.

Heavy metal	Pb (lead) Hg (mercury)	< 1 ppm < 0.03 ppm
specification	Cd (cadmium)	< 0.1 ppm

\* Analysed on regular basis.

Microbiological specification	Bacillus cereus	<100 CFU/g	Method: Sacco M10 (1)
	Coagulase positive staphylococci*	<10 CFU/g	Method: Sacco M11(2)
	Enterobacteriaceae	<10 CFU/g	Method: Sacco M2 (3)
	Escherichia coli	<1 CFU/g	Method: Sacco M27 (4)
	Listeria monocytogenes*	Not detected in 25 g	Method: Sacco M13 (5)
	Moulds & yeasts	<10 CFU/g	Method: Sacco M3 (6)
	Salmonella spp*	Not detected in 25 g	Method: Sacco M12 (7)

\* Analysed on regular basis. All analytical methods are available upon request. (1)ISO 7932; (2)ISO 6888-1-2; (3)ISO 21528-1-2; (4)ISO11866-1-2/IDF 170-1-2; (5)ISO 11290-1-2; (6)ISO 6611/IDF 94; (7)ISO 6785/IDF 93.

**GMO** The microbial strains are not genetically modified (GMO) in accordance with the European Directive 2001/18/EC. The strains are isolated from natural sources. The raw materials used are also GMO free in accordance with Regulation (EC) No. 1829/2003 and Regulation (EC) No. 1830/2003. Statement available upon request.

## Allergens The raw materials used are generally based on dairy ingredients. All materials are free of the following components and their derivates: peanut, tree nut, sesame, egg, fish, shellfish, mollusc, crustacean, sulphite, cereals containing gluten, celery, mustard, soy and lupine. Statement available upon request.

Safety information Material Safety Data Sheet available on www.saccosrl.it

Certificate Lot certificate available upon request.

**ISO** Sacco S.r.I. is UNI EN ISO 9001:2008 certified since 1998, ISO 22000:2005 and FSSC 22000 certified since 2014. Sacco cultures are generally Kosher approved except for surface ripening cultures.

- Service Please contact your distributor for guidance and instructions for your choice of culture and processing. Information about additional package sizes and sales units is also available upon request.
- Liability This information is based on our knowledge trustworthy and presented in good faith. No guarantee against patent infringement is implied or inferred.